

Extensions of endomorphisms of $C(X)$

J. F. Feinstein and T. J. Oliver

Abstract

For a compact space X we consider extending endomorphisms of the algebra $C(X)$ to be endomorphisms of Arens-Hoffman and Cole extensions of $C(X)$. Given a non-linear, monic polynomial $p \in C(X)[t]$, with $C(X)[t]/pC(X)[t]$ semi-simple, we show that if an endomorphism of $C(X)$ extends to the Arens-Hoffman extension with respect to p then it also extends to the simple Cole extension with respect to p . We show that the converse to this is false. For locally connected, metric X we characterize the algebraically closed $C(X)$, in terms of the extendability of endomorphisms to Arens-Hoffman and to simple Cole extensions.

1 Introduction

The problem of classifying the endomorphisms of commutative Banach algebras and their spectra has been extensively studied (see, for example, [5, 6, 10, 11, 12]). By contrast, the extendability problem for endomorphisms of commutative Banach algebras appears to be relatively untouched. In this note we concentrate on two different types of extension of the algebra $C(X)$ of continuous, complex-valued functions on X obtained by adjoining a root of some polynomial $p \in C(X)[t]$: the Arens-Hoffman extension and the ‘simple’ Cole extension. For a thorough treatment of these algebra extensions and how they relate to each other we refer the reader to [4] (see also [1, 2]). Throughout this paper, all Cole extensions discussed will be assumed to be simple in the above sense.

The question of whether or not an endomorphism of $C(X)$ extends to a given Cole extension may be viewed as purely topological. The question of whether or not an endomorphism of $C(X)$ extends to an Arens-Hoffman extension is more algebraic in nature. By asking these two questions simultaneously we gain insight into the relationships between the associated topological and algebraic conditions. We shall see, in the case when $C(X)[t]/pC(X)[t]$ is semi-simple, that if an endomorphism of $C(X)$ extends to the Arens-Hoffman extension with respect to p then it also extends to the Cole extension with respect to p . We shall also give an example to show that the converse is false. These results are, perhaps, surprising because the

Arens-Hoffman extension may be viewed as a sub-algebra of the Cole extension. Finally, for locally connected, compact metric spaces X , we shall characterize when $C(X)$ is algebraically closed in terms of the extendability of endomorphisms to Cole or Arens-Hoffman extensions.

2 Preliminary notation, definitions and results

Let W and X be topological spaces and A be a commutative algebra. We denote the ring of polynomials with coefficients in A by $A[t]$. The set of continuous functions from W to X is denoted by $C(W, X)$. The set of complex-valued, continuous functions on X is denoted by $C(X)$ which, when X is compact, is a commutative, unital, Banach algebra under pointwise addition and multiplication with the uniform norm $\|g\| := \sup\{|g(x)| : x \in X\}$. For $g \in C(X)$ we denote the zero set of g by $Z(g)$. For $q := q_0 + \dots + q_m t^m \in C(X)[t]$ and a homomorphism $\alpha : C(X) \rightarrow A$ we define $q^{(\alpha)} = \alpha(q_0) + \dots + \alpha(q_m)t^m \in A[t]$. For $\beta \in C(W, X)$ we define $\beta^* : C(X) \rightarrow C(W)$ by $\beta^*(g) = g \circ \beta$ ($g \in C(X)$).

In our terminology a *compact space* is a compact, Hausdorff topological space. For a compact space X and $x \in X$, \hat{x} denotes the evaluation character on $C(X)$ at x : $\hat{x}(g) = g(x)$. Given a monic polynomial $p \in C(X)[t]$ of degree at least 2 we define $X_p = \{(x, \lambda) \in X \times \mathbb{C} : p^{(\hat{x})}(\lambda) = 0\}$. We define $\pi_X : X_p \rightarrow X$ and $\pi_p : X_p \rightarrow \mathbb{C}$ to be the restrictions to X_p of the two coordinate projections. The discriminant $D_p \in C(X)$ of p is defined by $D_p(x) = \prod_{(i < j)} (\lambda_i - \lambda_j)^2$ where $p^{(\hat{x})} = (t - \lambda_1) \times \dots \times (t - \lambda_n)$. If $\text{int}(Z(D_p)) = \emptyset$ then D_p is not a zero-divisor in $C(X)$ and p is of minimal degree in $C(X)[t] \setminus \{0\}$ such that $p^{(\pi_X^*)}(\pi_p) = 0$. In this case, the Arens-Hoffman extension $C(X)[t]/pC(X)[t]$ of $C(X)$ with respect to p is semi-simple and is isomorphic to the algebra $C(X)_p := \{q^{(\pi_X^*)}(\pi_p) : q \in C(X)[t]\} \subseteq C(X_p)$ ([1]). Note that $C(X)_p$ is the sub-algebra of $C(X_p)$ generated by $\{\pi_p\} \cup \pi_X^*(C(X))$. In this setting, the Cole extension of $C(X)$ with respect to p is simply $C(X_p)$ and this is equal to the uniform closure of $C(X)_p$ ([2, 4]). For convenience, we shall describe monic polynomials p of degree at least two and with $\text{int}(Z(D_p)) = \emptyset$ as *admissible* polynomials.

Throughout this note, all algebras considered will be commutative, unital, complex algebras. If A is an algebra we denote the set of unital endomorphisms of A by $\text{End}(A)$. Let B be an algebra and $\gamma : A \rightarrow B$ be an injective homomorphism. Clearly $\gamma(A)$ is a sub-algebra of B , and on $\gamma(A)$ we may define an inverse to γ which we denote by γ^{-1} . For $S \in \text{End}(A)$ we see that $\gamma S \gamma^{-1} \in \text{End}(\gamma(A))$. For $T \in \text{End}(B)$ we see that $T|_{\gamma(A)} = \gamma S \gamma^{-1}$ if and only if $T\gamma = \gamma S$. In this case we say that S *extends to B via γ* , and T *is an extension of S via γ* .

For a compact space X , the map $\phi \mapsto \phi^*$ (with notation as above) is a bijection from $C(X, X)$ to $\text{End}(C(X))$. The inverse map is given by $T \mapsto \phi_T$

where (regarding X as a subset of $C(X)^*$ in the usual way) $\phi_T = T^*|_X$. In this setting we say that T is the endomorphism of $C(X)$ induced by ϕ_T and that ϕ_T is the self-map of X associated with T .

We now give two observations which the reader may find useful to check before proceeding.

- (i) Let X be a compact space, $p, q \in C(X)[t]$ be monic, and $f \in C(X_p)$. Then $q^{(\pi_X^*)}(f) = 0$ if and only if $(x, f(x, \lambda)) \in X_q$ for all $(x, \lambda) \in X_p$.
- (ii) Let X be a compact space, $p \in C(X)[t]$ be monic, and $T \in \text{End}(C(X))$. Then $(x, \lambda) \in X_{p(T)}$ if and only if $(\phi_T(x), \lambda) \in X_p$.

3 Extending endomorphisms

We begin our investigation with some lemmas concerning the extendability of endomorphisms to the two types of algebra extension under consideration.

Recall that we defined the term *admissible* for polynomials in the previous section.

Lemma 1: Let X be a compact space, let $p \in C(X)[t]$ be admissible, and let $T \in \text{End}(C(X))$.

- (i) There is an extension $T_1 \in \text{End}(C(X)_p)$ of T via π_X^* if and only if there exists $f \in C(X)_p$ with $(x, f(x, \lambda)) \in X_{p(T)}$ for all $(x, \lambda) \in X_p$.
- (ii) Given an f satisfying the condition in (i), we may find an extension T_1 such that $T_1(\pi_p) = f$.

Proof: (i) Suppose that such an extension T_1 exists. Then $0 = T_1(0) = T_1(p^{(\pi_X^*)}(\pi_p)) = p^{(\pi_X^* T)}(T_1(\pi_p))$. That is $p^{(\pi_X^* T)}$ has a root $T_1(\pi_p) \in C(X)_p$. Set $f := T_1(\pi_p)$. Clearly $p^{(\pi_X^* T)}(f) = 0$, which means that $(x, f(x, \lambda)) \in X_{p(T)}$ for all $(x, \lambda) \in X_p$.

Conversely, suppose that such an f exists. Setting $T_1(\pi_p) := f$ and extending as an algebra homomorphism gives us a well defined endomorphism T_1 . To check that T_1 is well defined is easy when we note that p is monic and of minimal degree in $C(X)[t] \setminus \{0\}$ such that $p^{(\pi_X^*)}(\pi_p) = 0$.

Statement (ii) follows from the last part of the argument above. \square

Lemma 2: Let X be a compact space, let $p \in C(X)[t]$ be admissible, and let $T \in \text{End}(C(X))$. Then the following three statements are equivalent.

- (i) There exists an extension $T_2 \in \text{End}(C(X_p))$ of T via π_X^* .
- (ii) There exists $\tilde{\phi} \in C(X_p, X_p)$ such that $\pi_X \circ \tilde{\phi} = \phi_T \circ \pi_X$.

(iii) There exists $f \in C(X_p)$ with $(x, f(x, \lambda)) \in X_{p^{(T)}}$ for all $(x, \lambda) \in X_p$.

Moreover, given such an f satisfying the condition in (iii), we may find an extension T_2 such that $T_2(\pi_p) = f$.

Proof: The equivalence of (i) and (ii) follows immediately from the correspondence between self-maps and endomorphisms discussed above.

To see that (i) implies (iii), suppose that such a T_2 exists. Then setting $f := T_2(\pi_p) \in C(X_p)$ gives $p^{(\pi_X^* T)}(f) = p^{(T_2 \pi_X^*)}(T_2(\pi_p)) = T_2(p^{(\pi_X^*)}(\pi_p)) = 0$, and so $(x, f(x, \lambda)) \in X_{p^{(T)}}$ for all $(x, \lambda) \in X_p$.

Finally we prove that (iii) implies (ii), and simultaneously prove the last part of the statement. Suppose that such an f exists as in (iii). Define $\tilde{\phi} \in C(X_p, X_p)$ by $\tilde{\phi}(x, \lambda) := (\phi_T(x), f(x, \lambda))$ for all $(x, \lambda) \in X_p$. Set $T_2 = (\tilde{\phi})^*$. Clearly (i) and (ii) hold and $T_2(\pi_p) = f$. \square

Comparing Lemmas 1 and 2 we immediately obtain the following corollary.

Corollary 3: Let X be a compact space, let $p \in C(X)[t]$ be admissible, and let $T \in \text{End}(C(X))$. If T may be extended to $C(X)_p$ via π_X^* then T may also be extended to $C(X_p)$ via π_X^* .

Another way to see this is to consider the operator norm of an extension of T to $C(X)_p$ when $C(X)_p$ is given the uniform norm rather than the Arens-Hoffman norm.

This result leads us to the following three questions.

Question 1: Do there exist a compact space X , admissible $p \in C(X)[t]$, and $T \in \text{End}(C(X))$ such that T extends to $C(X)_p$ via π_X^* but there exists an extension $T_2 \in \text{End}(C(X_p))$ of T via π_X^* with $T_2|_{C(X)_p} \notin \text{End}(C(X)_p)$?

Question 2: Do there exist a compact space X , admissible $p \in C(X)[t]$, and $T \in \text{End}(C(X))$ such that T extends to $C(X_p)$ via π_X^* but T does not extend to $C(X)_p$ via π_X^* ?

Question 3: Do there exist a compact space X , admissible $p \in C(X)[t]$, and $T \in \text{End}(C(X))$ such that T does not extend to $C(X_p)$ via π_X^* ?

In order to answer Question 1 we need the following lemma.

Lemma 4: Let X be a compact space, let $p \in C(X)[t]$ be admissible, $f \in C(X)_p$, and $Y \subseteq X$ be a continuum. If y_0 is an interior point of Y , $\lambda_1, \lambda_2 \in C(Y)$ are such that $\lambda_1(y) \neq \lambda_2(y)$ for all $y \in Y \setminus \{y_0\}$, and $(y, \lambda_1(y)), (y, \lambda_2(y)) \in X_p$ for all $y \in Y$ then $\lim_{y \rightarrow y_0} \left(\frac{f(y, \lambda_1(y)) - f(y, \lambda_2(y))}{\lambda_1(y) - \lambda_2(y)} \right)$

exists and is finite.

Proof: Choose $q := q_0 + \dots + q_m t^m \in C(X)[t]$ such that $q^{(\pi_X^*)}(\pi_p) = f$. For all $y \in Y \setminus \{y_0\}$ we have

$$\frac{f(y, \lambda_1(y)) - f(y, \lambda_2(y))}{\lambda_1(y) - \lambda_2(y)} = \sum_{k=1}^m h_k(y),$$

where $h_k(y) = q_k(y)(\lambda_1(y)^{k-1} + \lambda_1(y)^{k-2}\lambda_2(y) + \dots + \lambda_2(y)^{k-1})$. It follows immediately that the required limit exists and is finite. \square

We now answer Question 1 in the affirmative by way of an example.

Example 1: Let $X := [0, 1]$ and define $r \in C(X)$ by $r(x) := (3x - 1)(3x - 2)^2$ for all $x \in X$. The properties of r that will be useful to us are that $r(\frac{1}{3}) = r(\frac{2}{3}) = 0$, $r'(\frac{1}{3}) \neq 0$, and $r'(\frac{2}{3}) = 0$. Now set $p(t) := (t - r)(t + r) \in C(X)[t]$ and let T be the automorphism of $C(X)$ induced by the self-homeomorphism $\phi(x) = 1 - x$ of X .

Setting $f \in C(X_p)$ to be $f(x, \lambda) := r(1 - x)$ for $\lambda = r(x)$, and $f(x, \lambda) := -r(1 - x)$ for $\lambda = -r(x)$ we have that $(x, f(x, \lambda)) \in X_{p(T)}$ for all $(x, \lambda) \in X_p$. So, by Lemma 2, there exists an extension $T_2 \in \text{End}(C(X_p))$ of T via π_X^* with $T_2(\pi_p) = f$. We see that $f \notin C(X)_p$ otherwise we would have, by Lemma 4, that $\lim_{x \rightarrow \frac{2}{3}} \left(\frac{f(x, r(x)) - f(x, -r(x))}{r(x) - (-r(x))} \right) = \lim_{x \rightarrow \frac{2}{3}} \left(\frac{r(1-x)}{r(x)} \right)$ exists and is finite. However, the aforementioned properties of r show that this is not the case. Thus, by Lemma 1, this particular T_2 does not restrict to give an endomorphism of $C(X)_p$.

Setting $f \in C(X_p)$ to be $f(x, \lambda) := r(1 - x)$ for all $(x, \lambda) \in X_p$ we still have that $(x, f(x, \lambda)) \in X_{p(T)}$ for all $(x, \lambda) \in X_p$, but we also have that $f \in C(X)_p$, since $f = \pi_X^*(T(r))$. So, by Lemma 1, T extends to $C(X)_p$ via π_X^* . \square

In order to answer Question 2 we need another lemma.

Definition: Let $S \subset \mathbb{T} \times \mathbb{C}$, where $\mathbb{T} := \{\lambda \in \mathbb{C} : |\lambda| = 1\}$. We call S a k -strip with respect to μ if $k \in \mathbb{N}$ and $\mu : \mathbb{T} \rightarrow S$ is a homeomorphism such that $\mu(e^{i\theta}) \in \{e^{ik\theta}\} \times \mathbb{C}$ for all $e^{i\theta} \in \mathbb{T}$.

Lemma 5: Let $a, b \in \mathbb{N}$, A be an a -strip with respect to μ_A , and B be a b -strip with respect to μ_B . There exists $g \in C(A, B)$ with $g(\{e^{i\theta}\} \times \mathbb{C}) \cap A \subset \{e^{i\theta}\} \times \mathbb{C}$ for all $e^{i\theta} \in \mathbb{T}$ if and only if b divides a . In this case $g(A) = B$.

Proof: Suppose that such a g exists. Consider $\mu_B^{-1} \circ g \circ \mu_A \in C(\mathbb{T}, \mathbb{T})$. Due to the properties of μ_A, μ_B , and g we see that $\mu_B^{-1} \circ g \circ \mu_A(e^{i\theta}) = e^{i(\frac{a}{b}\theta + \theta_0)}$




Figure 2

The sets \mathbb{T}_p and $\mathbb{T}_{p(T)}$ are given by

$$\mathbb{T}_p = \{(e^{i\theta}, \lambda) : (\theta, \lambda) \in [0, 2\pi]_{p(\beta^*)}\},$$

$$\mathbb{T}_{p(T)} = \{(e^{i\theta}, \lambda) : (\theta, \lambda) \in [0, 2\pi]_{p(\beta^*T)}\}.$$

Thus each of the sets \mathbb{T}_p , $\mathbb{T}_{p(T)}$ are the union of a 2-strip with a 3-strip. Denote the 2-strip in \mathbb{T}_p by R_2 , the 3-strip in \mathbb{T}_p by R_3 , the 2-strip in $\mathbb{T}_{p(T)}$ by S_2 , and the 3-strip in $\mathbb{T}_{p(T)}$ by S_3 . In view of Lemma 5, setting $g(x, \lambda) := (x, f(x, \lambda))$ for all $(x, \lambda) \in \mathbb{T}_p$, we see that there is exactly one $f \in C(\mathbb{T}_p)$ such that $(x, f(x, \lambda)) \in \mathbb{T}_{p(T)}$ for all $(x, \lambda) \in \mathbb{T}_p$. This f is such that g drops R_2 onto S_2 , and R_3 onto S_3 . Now by a similar argument to the one used in Example 1, considering $\lim_{\theta \rightarrow \pi} \left(\frac{f(e^{i\theta}, \lambda_2(\theta)) - f(e^{i\theta}, \lambda_5(\theta))}{\lambda_2(\theta) - \lambda_5(\theta)} \right)$, we see that $f \notin C(\mathbb{T})_p$. So T extends to $C(\mathbb{T}_p)$ via $\pi_{\mathbb{T}}^*$ but T does not extend to $C(\mathbb{T})_p$ via $\pi_{\mathbb{T}}^*$. \square

Our third example shows that the answer Question 3 is also affirmative, and again we may take X to be \mathbb{T} .

Example 3: Choose $p \in C(\mathbb{T})[t]$ as in Example 2. Let $T \in \text{End}(C(\mathbb{T}))$ be associated with $\phi_T \in C(\mathbb{T}, \mathbb{T})$ where $\phi_T(e^{i\theta}) := e^{i(\theta+\pi)}$ for all $\theta \in \mathbb{R}$. Denote the 2-strip in \mathbb{T}_p by R_2 , and the 3-strip in \mathbb{T}_p by R_3 . Finally, denote the 2-strip in $\mathbb{T}_{p(T)}$ by S_2 , and the 3-strip in $\mathbb{T}_{p(T)}$ by S_3 .

Suppose, for a contradiction, that there exists $f \in C(\mathbb{T}_p)$ such that $(x, f(x, \lambda)) \in \mathbb{T}_{p(T)}$ for all $(x, \lambda) \in \mathbb{T}_p$. Define $g \in C(\mathbb{T}_p, \mathbb{T}_{p(T)})$ by $g(x, \lambda) :=$

$(x, f(x, \lambda))$ for all $(x, \lambda) \in \mathbb{T}_p$. We see, as in Example 2, that g must drop R_2 onto S_2 , and R_3 onto S_3 . Hence g is surjective. We note that $(\{-1\} \times \mathbb{C}) \cap \mathbb{T}_p$ contains exactly four elements but $(\{-1\} \times \mathbb{C}) \cap \mathbb{T}_{p(T)}$ contains exactly five elements, as $\phi_T(-1) \neq -1$. This is a contradiction since g is single valued. Hence no such $f \in C(\mathbb{T}_p)$ may exist. Thus, by Lemma 2, T does not extend to $C(\mathbb{T}_p)$ via $\pi_{\mathbb{T}}^*$. \square

Careful consideration of the possible structures for X_p and $X_{p(T)}$ shows that, when $X = \mathbb{T}$, no admissible polynomial of degree less than 5 can lead to a positive answer to Question 3. If, however, you move to higher-dimensional X the situation is different. The following simple example of this phenomenon was suggested to us by J. W. Barrett. Take $X = \mathbb{T} \times \mathbb{T}$, let ϕ be the self-homeomorphism of X given by $\phi(z, w) = (w, z)$ and let T be the automorphism of $C(X)$ induced by ϕ . Now let F be the projection onto the first coordinate and let p be the quadratic polynomial $t^2 - F$ (which is clearly admissible, and indeed has invertible discriminant). Then it is easy to see that T has no extension to $C(X_p)$ via π_X^* .

4 Characterizing the algebraically closed $C(X)$

An algebra A is said to be *algebraically closed* if every non-constant, monic polynomial with coefficients in A has a root in A . In [3], Countryman characterized the first-countable compact spaces for which $C(X)$ is algebraically closed in terms of certain hereditary topological conditions on X which we describe below. He also showed that, for such X , $C(X)$ is algebraically closed if and only if $C(X) = \{g^2 : g \in C(X)\}$, and gave an example to show that this latter equivalence fails if X is not assumed to be first-countable.

For locally connected compact spaces, a different approach was taken by Hatori, Miura and Nijima ([7, 8]). They showed that, for such X , $C(X)$ is algebraically closed if and only if the covering dimension of X is at most 1 and $H^1(X, \mathbb{Z})$ is trivial, where $H^1(X, \mathbb{Z})$ is the first Čech cohomology group with integer coefficients. (It is perhaps worth noting that the combination of the latter two conditions is also equivalent to the condition that $\exp(C(X))$ be dense in $C(X)$.) For locally connected compact spaces it is again sufficient to be able to factorize the quadratic monic polynomials: $C(X)$ is algebraically closed if and only if $C(X) = \{g^2 : g \in C(X)\}$.

We now investigate the connection between $C(X)$ being algebraically closed and the extendability of endomorphisms to Arens-Hoffman extensions and Cole extensions. We begin with a lemma which immediately shows the connection in one direction.

Lemma 6: Let X be a compact space, $p \in C(X)[t]$ be admissible, and $T \in \text{End}(C(X))$. If $p^{(T)}(r) = 0$ for some $r \in C(X)$ then T extends to $C(X)_p$

via π_X^* .

Proof: Clearly $(x, r(x)) \in X_{p(x)}$ for all $(x, \lambda) \in X_p$. Set $f := \pi_X^*(r) \in C(X)_p$ and apply Lemma 1. \square

We now introduce the topological conditions used by Countryman to characterize when $C(X)$ is algebraically closed.

Definition: Let X be a compact space. We say that X is *hereditarily unicoherent* if every pair of continua in X have connected or empty intersection. We say that X is *not almost locally connected due to* $((C_n), (x_n), (y_n))$ if (C_n) is a sequence of pairwise disjoint continua in X which are open in $\overline{\bigcup_{n \in \mathbb{N}} C_n}$ such that $x_n, y_n \in C_n$ for all n , $x_n \rightarrow x_0 \in X$, and $y_n \rightarrow y_0 \in X \setminus \{x_0\}$. If no such $((C_n), (x_n), (y_n))$ exist then X is *almost locally connected*.

We may now state Countryman's characterization [3]: for a first-countable compact space X , $C(X)$ is algebraically closed if and only if X is hereditarily unicoherent and almost locally connected.

For the rest of this paper we take an arc to mean a homeomorphic copy of $[0, 1]$.

Lemma 7: Every almost locally connected, metric continuum is arc connected.

Proof: In [3] it is shown that every sequentially compact, almost locally connected Hausdorff continuum is locally connected. Thus every almost locally connected, metric continuum is locally connected. Every locally connected, metric continuum is arc connected, by Theorem 3.15 of [9]. Therefore every almost locally connected, metric continuum is arc connected. \square

Lemma 8: Let X be a locally connected, compact metric space. Then $C(X)$ is algebraically closed if and only if X does not contain a homeomorphic copy of \mathbb{T} .

Proof: We use Countryman's characterization, described above. It is clear that if X contains a homeomorphic copy of \mathbb{T} then X is not hereditarily unicoherent, and so $C(X)$ is not algebraically closed. Conversely, suppose that $C(X)$ is not algebraically closed.

Consider first the case where X is almost locally connected but not hereditarily unicoherent. We may choose continua $M, N \subset X$ such that $M \cap N = A \cup B$ where A and B are disjoint, non-empty compact subsets of X . Since M and N must also be almost locally connected we know, by Lemma 7, that M and N are arc connected. It is now easy to see that $M \cup N$

contains a homeomorphic copy of \mathbb{T} .

The remaining case is where X is not almost locally connected due to some $((C_n), (x_n), (y_n))$. Let X_0 be the component of X containing x_0 and y_0 . As X is locally connected we see that X_0 is open. So there is an $n_0 \in \mathbb{N}$ such that $C_n \subset X_0$ for $n \geq n_0$. Each C_n is open in $\overline{\bigcup_{n \in \mathbb{N}} C_n}$, and X_0 is locally arc connected, by Lemma 3.29 of [9]. So for each $n \geq n_0$, and $x \in C_n$ there is a $d_x > 0$ such that $B_{d_x}(x)$ is arc connected, and $B_{2d_x}(x) \cap C_m = \emptyset$ for all $m \neq n$. As C_n is compact we may cover C_n with finitely many such $B_{d_x}(x)$. Hence we may assume that the C_n are arcs from x_n to y_n . We now have $x_n \rightarrow x_0$, $y_n \rightarrow y_0$, each C_n is an arc from x_n to y_n , and there exist disjoint, arc connected neighbourhoods of x_0 and y_0 . It follows easily that X_0 contains a homeomorphic copy of \mathbb{T} . \square

Theorem 9: Let X be a locally connected, compact metric space. Then the following are equivalent.

- (i) $C(X)$ is algebraically closed.
- (ii) For all closed $Y \subseteq X$, admissible $p \in C(Y)[t]$, and $T \in \text{End}(C(Y))$ we have that T extends to $C(Y)_p$ via π_Y^* .
- (iii) For all closed $Y \subseteq X$, admissible $p \in C(Y)[t]$, and $T \in \text{End}(C(Y))$ we have that T extends to $C(Y_p)$ via π_Y^* .

Proof: Suppose, first, that (i) does not hold. Then, by Lemma 8, there is a $Y \subseteq X$ and a homeomorphism $h : Y \rightarrow \mathbb{T}$. Choose $p \in C(\mathbb{T})[t]$ and $T \in \text{End}(C(\mathbb{T}))$ as in Example 3. Let $T_h \in \text{End}(C(Y))$ be associated with $h^{-1} \circ \phi_T \circ h \in C(Y, Y)$. Since T does not extend to $C(\mathbb{T}_p)$ via $\pi_{\mathbb{T}}^*$ we have that T_h does not extend to $C(Y_{p^{(h^*)}})$ via π_Y^* . Hence (iii) does not hold, and so, by Corollary 3 (ii) does not hold.

Suppose, now, that (i) holds. Then for all closed $Y \subseteq X$ we have that $C(Y)$ is algebraically closed. Thus for admissible $p \in C(Y)[t]$ and $T \in \text{End}(C(Y))$ we have that $p^{(T)}$ has a root in $C(Y)$. Lemma 6 now shows that (ii) holds and hence, by Corollary 3, (iii) holds. \square

We conclude with a conjecture.

Conjecture: Let X be a first countable, compact space. Then the following are equivalent.

- (i) $C(X)$ is algebraically closed.
- (ii) For all admissible $p \in C(X)[t]$ and $T \in \text{End}(C(X))$ we have that T extends to $C(X)_p$ via π_X^* .

- (iii) For all admissible $p \in C(X)[t]$ and $T \in \text{End}(C(X))$ we have that T extends to $C(X_p)$ via π_X^* .

References

- [1] R. Arens and K. Hoffman, *Algebraic extension of normed algebras*, Proc. Am. Math. Soc. **7** (1956), 203–210.
- [2] B. J. Cole, *One-point parts and the peak point conjecture*, Ph.D. Thesis, Yale University, 1968.
- [3] R. S. Countryman, *On the characterization of compact Hausdorff X for which $C(X)$ is algebraically closed*, Pacific J. Math. **20** (1967), 433–448.
- [4] T. W. Dawson, *A survey of algebraic extensions of commutative, unital normed algebras*, Contemp. Math. **328** (2003), 157–170.
- [5] J. F. Feinstein and H. Kamowitz, *Endomorphisms of algebras of infinitely differentiable functions on compact plane sets*, J. Funct. Anal. **173** (2000), 61–73.
- [6] P. Galindo, T. W. Gamelin and M. Lindström, *Composition Operators on Uniform Algebras and the Pseudohyperbolic Metric*, J. Korean Math. Soc. **41** (2004), 1–20.
- [7] O. Hatori and T. Miura, *On a characterization of the maximal ideal spaces of commutative C^* -algebras in which every element is the square of another*, Proc. Amer. Math. Soc. **128** (2000), 1185–1189.
- [8] T. Miura and K. Nijjima, *On a characterization of the maximal ideal spaces of algebraically closed commutative C^* -algebras*, Proc. Amer. Math. Soc. **131** (2003), 2869–2876.
- [9] J. G. Hocking and G. S. Young, *Topology*, Dover Publications Inc., New York, 1961.
- [10] H. Kamowitz, *Compact endomorphisms of Banach algebras*, Pacific J. Math. **89** (1980), 313–325.
- [11] H. Kamowitz and S. Scheinberg, *Some properties of endomorphisms of Lipschitz algebras*, Studia Math. **96** (1990), 255–261.
- [12] U. Klein, *Kompakte multiplikative Operatoren auf uniformen Algebren*, Mitt. Math. Sem. Giessen **232** (1997).

School of Mathematical Sciences
University of Nottingham
Nottingham NG7 2RD, England
Joel.Feinstein@nottingham.ac.uk
Thomas.Oliver@maths.nottingham.ac.uk

2000 Mathematics Subject Classification: 46J10, 47B48

The second author would like to thank the EPSRC for providing support for this research